

Tunneling time scale of under-the-barrier forerunners

Gastón García-Calderón^{1,*} and Jorge Villavicencio^{2,†}

¹*Instituto de Física, Universidad Nacional Autónoma de México,
Apartado Postal 20 364, 01000 México, Distrito Federal, México*

²*Facultad de Ciencias, Universidad Autónoma de Baja California,
Apartado Postal 1880, 22800 Ensenada, Baja California, México*

(Dated: February 9, 2008)

Time-dependent analytical solutions to Schrödinger's equation with quantum shutter initial conditions are used to investigate the issue of the tunneling time of forerunners in rectangular potential barriers. By using a time-frequency analysis, we find the existence of a regime characterized by the opacity of the barrier, where the maximum peak of a forerunner measured at the barrier transmission edge $x = L$ corresponds to a genuine tunneling process. The corresponding time scale represents the tunneling time of the forerunner through the classically forbidden region.

PACS numbers: 03.65Bz, 03.65Ca.

Tunneling, that describes the possibility that a particle traverses through a classically forbidden region, is one of the paradigms of quantum mechanics. In the energy domain one solves Schrödinger's equation at a fixed energy E to obtain the probability of transmission through a barrier region, a subject discussed in every quantum mechanics textbook. In the time domain, however, there are still aspects open to scrutiny. A problem that has remained controversial over the years is the tunneling time problem, that may be stated by the question : How long does it take to a particle to traverse a classically forbidden region? Different authors have proposed and defended different views in answering the above question[1]. Some authors have argued that it might not be a unique definition of the tunneling time since different arrangements for the tunneling process may lead to different relevant time scales. An interesting idea that arises from the above considerations is that the relevance of the different tunneling time definitions may depend on the context where they may become physically significant quantities[2].

In recent work we have investigated the effect of the transient solutions to the time-dependent Schrödinger's equation for cutoff wave initial conditions (quantum shutter) on the tunneling process[3, 4, 5]. In particular for tunneling through a barrier we found that just across the tunneling barrier, the probability density as a function of time may exhibit a sharp peak maximum that we called *time domain resonance*. The peak value t_{max} of that forerunner represents the largest probability of finding the particle at the barrier width L . We found also that the behavior of t_{max} as a function of the barrier width L exhibits for small values of L a *basin* region, followed by a region where t_{max} grows linearly as L increases[4]. More recently, in collaboration with Delgado and Muga[2], we have investigated the time scale for forerunners preceeding the main tunneling signal of a wave created by a source with a sharp onset[6, 7] and by a quantum shutter[4] for systems with very large or infinite barrier width. For the particular case of opaque finite

barriers [2] we found a basin regime occurring at fixed positions $0 < x \ll \kappa_0^{-1}$ along the internal region, where forerunners are dominated by under-the-barrier frequency components. That is, in opaque barrier systems the tunneling forerunners are observed at distances of the order or smaller than the penetration length κ_0^{-1} , where $\kappa_0 = [2m(V - E)]^{1/2}/\hbar$, with E and V corresponding to the incidence energy and the barrier height, respectively. However, at distances $\kappa_0^{-1} < x \leq L$ the non-tunneling components eventually dominate the time evolution process, and hence it is not possible to speak of a genuine tunneling time scale of forerunners at $x = L$. We believe that in order to gain more insight on the properties of transient tunneling structures and their corresponding time scales, the existence of tunneling forerunners at the barrier edge $x = L$ needs to be investigated.

The aim of this work is to show the existence of a regime along the *basin* region, where the maximum t_{max} of the *time domain resonance* measured at the barrier width L corresponds to under-the-barrier tunneling and consequently provides a genuine time scale for tunneling.

Our approach to the tunneling time problem is based on a model that deals with an explicit solution[3] of time-dependent Schrödinger equation for an arbitrary potential $V(x)$ ($0 \leq x \leq L$) that vanishes outside the internal region. We consider the problem of the time evolution of a cutoff plane wave $\Psi(x, k; t = 0) = \Theta(-x)(e^{ikx} - e^{-ikx})$ following the instantaneous opening at $t = 0$ of a quantum shutter at $x = 0$. Along the tunneling region the solution reads,

$$\Psi^i(x, k, t) = \Phi_k(x)M(y_k) - \Phi_{-k}(x)M(y_{-k}) - \sum_{n=-\infty}^{\infty} \Phi_n(x)M(y_{k_n}), \quad (0 \leq x \leq L) \quad (1)$$

where the $\Phi_{\pm k}(x)$'s refer to the stationary solutions of the problem and $\Phi_n(x) = 2iku_n(0)u_n(x)/(k^2 - k_n^2)$. Similarly, the solution $\Psi^e(x, k; t)$ for the external or transmitted region ($x \geq L$), is given by [4],

$$\Psi^e(x, k; t) = T_k M(y_k) - T_{-k} M(y_{-k}) - i \sum_{n=-\infty}^{\infty} T_n M(y_{k_n}), \quad (x \geq L) \quad (2)$$

where the $T_{\pm k}$'s refer to the transmission amplitudes, and the factor $T_n = 2ik u_n(0) u_n(L) \exp(-ik_n L) / (k^2 - k_n^2)$. In Eqs. (1) and (2) the coefficients Φ_n and T_n are given in terms of the resonant eigenfunctions $\{u_n(x)\}$ with complex energy eigenvalues $E_n = \hbar^2 k_n^2 / 2m$, with $k_n = a_n - ib_n$ ($a_n, b_n > 0$). The resonant sums in Eqs. (1) and (2) run over the full set of complex poles $\{k_n\}$. The M 's are defined as [3],

$$M(y_q) = \frac{1}{2} e^{imx^2/2\hbar t} w(iy_q) \quad (3)$$

where w is the complex error function [8] defined as $w(z) = \exp(-z^2) \text{erfc}(-iz)$, with arguments $y_q(x, t) = e^{-i\pi/4} (m/2\hbar t)^{1/2} [x - \hbar q t / m]$, where $q = \pm k, k_{\pm n}$.

We shall explore the issue of the tunneling time scale in typical one dimensional potential barriers [9] of height V and thickness L , defined along the interval $0 \leq x \leq L$. Let us analyze the behavior of the maximum of the *time domain resonance*, t_{max} , as a function of the potential barrier width, L . Let us recall that this transient structure corresponds to the first maximum of the probability density $|\Psi^e|^2$, measured at the fixed position, $x = L$. This can be appreciated in Fig. 1 where we plot the L -dependence of t_{max} (full dot), corresponding to potential barrier systems with parameters: $V = 0.3$ eV, incidence energy $E = \hbar^2 k^2 / 2m = 0.001$ eV, and effective mass for the electron $m = 0.067m_e$. Here we can clearly observe a *basin* corresponding to a range of values of the barrier width. We can also appreciate that if L is further increased, t_p grows linearly with L . Such a linear regime occurs at large barrier widths. We have argued [4] that in this case, the tunneling process is inhibited and that the particle goes mainly over the barrier. We have recently suggested [4] that for small values of the barrier width L , the *basin* exhibited by t_{max} , is a result of a subtle interplay between tunneling and top-barrier resonant processes. In what follows we shall investigate under what conditions the time scales associated to the *basin*, are in fact related to a tunneling process. We begin our analysis by choosing the case $L = 4.0$ nm depicted in Fig. 1, which is located around the minimum of the *basin*. In Fig. 2 we plot the normalized probability density $|\Psi^e|^2$ (solid line), normalized to the transmission coefficient $|T_k|^2$, as a function of time t for a fixed value of position $x = L$. We can clearly appreciate a *time domain resonance* peaked at the value $t_{max} = 5.17$ fs. In order to determine if this structure is related to a genuine tunneling process we analyze the frequency content of $|\Psi^e|^2$.




FIG. 1: Maximum of the *time domain resonance* t_{max} (full dot) as a function of the barrier width L , for an incidence energy $E = .001$ eV. In this case the barrier height is $V = 0.3$ eV. See text.




FIG. 2: Time evolution of $|\Psi^e|^2$ (solid line), at $x = L = 4.0$ nm. A full square indicates the position of the maximum of the *time-domain resonance*, at $t_{max} = 5.17$ fs.

We compute the local average frequency ω_{av} [6, 10],

$$\omega_{av} = -\text{Im} \left[\frac{1}{\Psi^s} \frac{d}{dt} \Psi^s \right], \quad (4)$$

and the instantaneous bandwidth σ [10],

$$\sigma = \left| \text{Re} \left[\frac{1}{\Psi^s} \frac{d}{dt} \Psi^s \right] \right|, \quad (5)$$

where $s = i, e$ according to consider, respectively, the internal or external solutions.

In Fig. 3 we plot the relative average local frequency (relative frequency for short) ω_{av}/ω_V , where $\omega_V = V/\hbar$ is the cut-off frequency, along the relevant time interval,




FIG. 3: Relative average local frequency ω_{av}/ω_V (solid line) for the case depicted in Fig. 2. The cutoff-frequency $\omega_V/\omega_V = 1$ (dashed line) is included for comparison. In the inset we plot the instantaneous bandwidth σ of the spectrogram depicted in the main graph. Notice that the frequency deviations from t_{max} are exactly zero, *i.e.*, $\sigma(t_{max}) = 0$. In all cases a full square indicates the position of t_{max} .

discussed in Fig. 2. We can appreciate that in the vicinity of the maximum of the *time-domain resonance*, t_{max} , the probability density is composed entirely by under-the-barrier frequency components *i.e.* $\omega_{av}/\omega_V < 1$. This also occurs at the exact value t_{max} , also indicated in the figure by a solid square. In the inset of Fig. 3 we plot the instantaneous bandwidth σ of the spectrogram. Notice the absence of a frequency dispersion around the maximum t_{max} , *i.e.*, $\sigma(t_{max}) = 0$. The above result indicates that in this case the peak of the *time domain resonance*, and the values close to it, refer to a genuine tunneling event. In Fig. 4 we plot the relative frequency ω_{av}/ω_V associated to different values of the maximum t_{max} of the *time-domain resonance*, measured at different positions along both the internal and external regions of a potential barrier with parameters: $V = 0.3$ eV, $L = 4.13$ nm, and $\alpha = 3.0$. In this case we choose the following values of the incidence energy: $E = 0.001$ eV (solid dot), and $E = 0.01$ eV (hollow dot). In the inset of Fig. 4 we show, for the particular case of $E = 0.01$ eV, the values of t_{max} (solid square) at the different values of position considered in the main graph. As can be clearly appreciated in that figure, the tunneling process along the whole internal region is governed by under-the-barrier-frequency components, *i.e.*, $\omega_{av}/\omega_V < 1$. The above results should be contrasted with those obtained for very large or infinite barrier widths recently studied in Ref. [2], where under-the-barrier tunneling components arise only at distances smaller than a characteristic length. Here we have found a particular combination of potential parameters characterizing a genuine tunneling process. The existence of under-the-barrier tunneling forerunners is not only re-




FIG. 4: Relative frequency ω_{av}/ω_V of the maximum of the *time-domain resonance*, as a function of position, measured in units of the barrier width L . The parameters are given in the text. Two incidence energies are considered: $E = 0.001$ eV (solid dot), and $E = 0.01$ eV (hollow dot). In both cases, the relative frequency ω_{av}/ω_V along the internal region is below the cutoff-frequency $\omega_V/\omega_V = 1$ (dashed line). The behavior of t_{max} as a function of position is illustrated in the inset for the case with $E = 0.01$ eV. The position of the barrier edge, $x = L$, is indicated by a dotted line in both figures.




FIG. 5: Relative frequency ω_{av}/ω_V measured at the barrier edge $x = L$, as a function of the opacity α . Here we considered a barrier height $V = 0.3$ eV, and the parameters: $u = 300$ (solid dot), $u = 10$ (solid triangle), and $u = 5$ (solid square). Note that for values of the opacity smaller than $\alpha \simeq 3.3$, the relative frequencies for all values of u are below the cutoff-frequency $\omega_V/\omega_V = 1$ (dashed line). See text.

stricted to values of the position along the internal region of the potential. In fact, we can see in Fig. 4, that for distances up to $x \simeq 2L$ along the external region, we can still observe frequency components below the cutoff-frequency, ω_V . As x/L increases further, $\omega_{av}/\omega_V > 1$.

This behavior follows because the corresponding values of t_{max} increase linearly, indicating the prevalence of non-tunneling components (see the inset in Fig. 4).

We have found that the existence of tunneling forerunners at the barrier edge $x = L$ may be described more generally by referring to the opacity α of the system, defined as,

$$\alpha = \frac{[2mV]^{1/2}}{\hbar} L, \quad (6)$$

and by the dimensionless parameter u , the ratio between the potential barrier height and the incidence energy,

$$u = \frac{V}{E}. \quad (7)$$

There is a regime, that we shall refer to as *tunneling regime*, characterized by the opacity α and the parameter u , where the relative frequencies associated to the *time domain resonance* are below the cutoff-frequency ω_V . In order to show this, in Fig. 5 we plot the relative frequency ω_{av}/ω_V as function of the opacity α , for three different values of the parameter u , $u = 5$ (solid dot), $u = 10$ (solid triangle), and $u = 300$ (solid square). Although in this case we have chosen a value of $V = 0.3$ eV, we have checked that the relative frequency ω_{av}/ω_V , exhibits a striking regularity when plotted as a function of α . For a given value of α all the systems with the same parameter u , share the same curve. That is, all systems characterized by the same parameters α and u yield the same relative frequency, ω_{av}/ω_V . In view of this observed regularity, we can completely characterize the regime associated with under-the-barrier frequency components. Although in Fig. 5 we have considered values of the parameter u such that $5 \leq u \leq 300$, the case corresponding to very large values of u ($u \rightarrow \infty$), almost overlaps with the case $u = 300$. Thus, for very large values of u there is an upper-bound for the opacity $\alpha_u \simeq 3.3$. This result, and the fact that the lower-bound for the opacity [11] is given by $\alpha_c = 2.065$, recently reported in Ref. [5], allows to characterize an opacity “window” given by $\alpha_c \leq \alpha \leq \alpha_u$ where the relative frequencies are always below the cutoff-frequency $\omega_V/\omega_V = 1$, irrespective of the value of the parameter u , namely of the value of the incidence energy. Note that the values of α within the opacity “window” may be obtained using typical parameters of semiconductor heterostructures[9]. From the figure one can appreciate the existence of another regime of α ’s where the existence of under-the-barrier processes associated to the *time-domain resonance* may depend on the value of the parameter u .

It is of interest to remark that the time scale given by the peak maximum t_{max} of the forerunner at the barrier edge $x = L$, i.e., the *time domain resonance*, differs from both the semi-classical Büttiker-Landauer and Büttiker traversal times, which exhibit a linear dependence with

L , and also differs from the phase-time[4, 5]. In Ref. [5] we have discussed an unexpected relationship between the delay time and the existence of *time domain resonances*.

To conclude we remark that the analytical solution to the time-dependent Schrödinger’s equation with quantum shutter initial conditions applies in general to arbitrary potentials provided they vanish beyond a distance and can also be extended to deal with cutoff pulses as discussed in Ref. [5]. Our results refer to novel transient effects in time-dependent tunneling. To test our results experimentally would require to consider the detection of tunneling particles in time domain at distances close to the interaction region.

The authors thank J. G. Muga for a critical reading of the manuscript and acknowledge financial support of DGAPA-UNAM under grant No. IN101301.

* **Senior Associate ICTP**; Electronic address: gaston@fisica.unam.mx

† Electronic address: villavics@uabc.mx

- [1] See for example: E. H. Hauge and J. A. Støvneng, Rev. Mod. Phys. **61**, 917 (1989); R. Landauer and Th. Martin, Rev. Mod. Phys. **66**, 217 (1994); P. Ghose, *Testing Quantum Mechanics on New Ground* (Cambridge University Press, Cambridge, 1999), Chapter 10; *Time in Quantum Mechanics*, J. G. Muga, R. Sala, I. L. Egusquiza (eds.) (Springer-Verlag, Berlin, 2002).
- [2] G. García Calderón, J. Villavicencio, F. Delgado, and J. G. Muga, Preprint arXiv: quant-ph 0206020 (2002). To be published in Phys. Rev. A.
- [3] G. García-Calderón, A. Rubio, Phys. Rev. A **55**, 3361 (1997).
- [4] G. García-Calderón and Jorge Villavicencio, Phys. Rev. A **64**, 012107 (2001).
- [5] G. García-Calderón and Jorge Villavicencio, Preprint arXiv: quant-ph 0210008 (2002), to be published in Phys. Rev. A.
- [6] J. G. Muga and M. Büttiker, Phys. Rev. A **62**, 023808 (2000).
- [7] J. Villavicencio, R. Romo, and S. Sosa y Silva, Preprint ArXiv: quant-ph 0210010 (2002). To be published in Phys. Rev. A.
- [8] *Handbook of Mathematical Functions*, edited by M. Abramowitz and I. A. Stegun (Dover, New York 1965), p. 297.
- [9] E. E. Mendez, in *Physics and Applications of Quantum Wells and Superlattices*, edited by E. E. Mendez and K. Von Klitzing (Plenum, New York, 1987) p. 159; D. K. Ferry and S. M. Goodnick, *Transport in Nanostructures*, Cambridge University Press, United Kingdom, 1997 (pp. 91-201).
- [10] L. Cohen, *Time-Frequency analysis* (Prentice Hall, New Jersey, 1995).
- [11] As discussed in Ref. [5], the lower-bound $\alpha_c = 2.0653$ correspond to a critical opacity, such that for values of $\alpha > \alpha_c$, *time-domain resonances* may exist.